



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

PROBLEMS FOR SOLUTION.

ALGEBRA.

253. Proposed by R. D. CARMICHAEL, Hartselle, Ala.

Prove that $x^5 + ax + b = 0$ is solvable by radicals if $b = ma$, m being the negative of half the sum of any two roots of the original equation. Exhibit the solution.

254. Proposed by R. D. CARMICHAEL, Hartselle, Ala.

Sum to infinity the series $\frac{n^2}{(16n^2 - 1)^2}$ beginning with $n = 1$.

255. Proposed by O. E. GLENN, Ph. D., Springfield, Mo.

Let f be the binary cubic $a_0x_1^3 + 3a_1x_1^2x_2 + 3a_2x_1x_1^2 + a_3x_2^3$, $\Delta = (f, f)_2$ the covariant, the second transvectant of f over itself, and $R = 2[4(a_0a_2 - a_1^2) \times (a_1a_3 - a_2^2) - (a_0a_3 - a_1a_2)^2] = (\Delta, \Delta)_2$ the second transvectant of Δ over itself. Then if $\Delta_{\kappa\lambda}$ is the Δ covariant for the cubic pencil $\kappa f + \lambda Q$, Q being the first transvectant of f over Δ we have $\Delta_{\kappa\lambda} = (\kappa^2 - \frac{1}{2}\lambda^2 R) \Delta$.

CALCULUS.

900. Proposed by PROFESSOR B. F. FINKEL, A. M., 4038 Locust Street, Philadelphia, Pa.

Prove that, if the differential equation $cydx - (y + a + bx)dy - nx(xdy - ydx) = 0$, be transformed into an equation between u and x by the substitution $u(y + a + bx + nx^2) = y(c + nx)$, then the variables are separable; and reduce the equation to the form $dv/\phi(v) = dx/\phi(x)$ by the further substitution $v = au + \beta$, a and β being suitably determined. *Euler*. [Forsyth's *Differential Equations*, p. 48, Ex. 4.]

DIOPHANTINE ANALYSIS.

132. Proposed by O. E. GLENN, Ph. D., Springfield, Mo.

Disregarding the order of λ, μ, ν , how many sets of solutions has the congruence $\lambda + \mu + \nu \equiv 0 \pmod{p-1}$ (p prime)?

GEOMETRY.

280. Proposed by WILLIAM HOOVER, Ph. D., Athens, Ohio.

On any diameter of a given ellipse is taken a point such that the tangents from it intercept on the tangent at one end of the diameter a length equal to the diameter; the ellipse being $a^2y^2 + b^2x^2 - a^2b^2 = 0$. Prove that the locus of the point is $\left(\frac{x^2}{a^2} - \frac{y^2}{b^2}\right)^2 = \left(\frac{a^2 + b^2}{a^2 - b^2}\right)^2 \left(\frac{x^2}{a^2} + \frac{y^2}{b^2}\right)$.